

Supersymmetric Extension of the Minimal Dark Matter Model

Xue Chang¹, Chun Liu¹, Feng-Cai Ma², Shuo Yang^{3,4}

¹ *State Key Laboratory of Theoretical Physics,
Institute of Theoretical Physics, Chinese Academy of Sciences,
P.O. Box 2735, Beijing 100190, China*

² *Physics Department, Liaoning University, Shenyang, 110036, P.R. China*

³ *Physics Department, Dalian University, Dalian, 116622, P.R. China*

⁴ *Center for High-Energy Physics, Peking University, Beijing, 100871, P.R. China**

Abstract

The minimal dark matter model is given a supersymmetric extension. A super $SU(2)_L$ quintuplet is introduced with its fermionic neutral component still being the dark matter, the dark matter particle mass is about 19.7 GeV. Mass splitting among the quintuplet due to supersymmetry particles is found to be negligibly small compared to the electroweak corrections. Other properties of this supersymmetry model are studied, it has the solutions to the PAMELA and Fermi-LAT anomaly, the predictions in higher energies need further experimental data to verify.

PACS numbers: 95.35.+d, 12.60.-i, 14.80.-j

*Electronic address: chxue@itp.ac.cn, liuc@mail.itp.ac.cn, fcma@lnu.edu.ca, yangshuo@dlu.edu.cn

I. INTRODUCTION

It has been known from many astrophysical measurements that the universe contains enormous of invisible, non-baryonic dark matter (DM) which is not included in the Standard Model (SM). Among various hypotheses for the nature of the DM, that of weakly interacting massive particles (WIMPs) is very attractive. Focussing on the DM problem, one can explore a simple WIMP model, that is the minimal dark matter model (MDM) [1, 2]: adding to the SM a single matter X without introducing any additional discrete symmetry, and X is in a high dimensional representation of the usual SM $SU(2)_L \times U(1)_Y$ electroweak (EW) symmetry. The stability of the DM candidate is guaranteed by the SM gauge symmetry and by the renormalizability. The minimality of the model lies in the fact that the new physics is determined by only one parameter, namely the mass M of the X multiplet. Therefore, the MDM is remarkably predictive. There are some extensions to the MDM [3].

As far as the particle physics is concerned, the SM provides a successfully description of presently known phenomena. However it will have to be extended to describe physics at higher energies. That often results in the gauge hierarchy problem. Supersymmetry (SUSY) offers a solution to this problem [4].

In this work, we make SUSY extension to the MDM. Note that in the so-called minimal SUSY extension of the SM (MSSM), the DM candidate, that is the lightest SUSY particle, is there only after introducing an extra discrete symmetry by hand, which is the R -parity. In our SUSY MDM (SMDM), instead, we still follow the logic of MDM, the existence of the DM lies in the fact that the DM is in a high dimensional representation of the SM gauge group without using discrete symmetries.

In section II, the SMDM is constructed. In section III, mass splitting of the X multiplet, the DM relic density, direct and indirect detection signatures of the SMDM are calculated. In section IV, the conclusion is made. In the Appendix, we give basic facts about the representation of the $SU(2)$ group.

II. SMDM

The SMDM is simply constructed by supersymmetric extension to the MDM. The particle content is, in addition to that of the MSSM, the fermionic $SU(2)_L$ 5-plet X of the MDM and

its superpartner which a complex scalar 5-plet \tilde{X} . The charged components X^Q are slightly heavier than the neutral one X^0 due to quantum corrections [1, 2]; and the superpartner \tilde{X} because of its soft mass M_{soft} , is also heavier than X . Both X^Q and \tilde{X} will decay into X^0 . The relic particle in the SMDM is still X^0 as in the MDM. The new parameters are M and M_{soft} , the model is still predictive. As for the Lagrangian, in addition to that of the MSSM, we have

$$\begin{aligned}\mathcal{L}_{SMDM} = & \frac{i}{2}(X^{\dagger i}\bar{\sigma}^\mu\partial_\mu X_i + \bar{X}^{\dagger i}\bar{\sigma}^\mu\partial_\mu \bar{X}_i) - \frac{i}{2}g_2 A_a^\mu(X^{\dagger i}\bar{\sigma}_\mu(T^a)_i{}^j X_j - \bar{X}^{\dagger i}\bar{\sigma}_\mu(T^a)_j{}^i \bar{X}_j) \\ & - \frac{\sqrt{2}}{2}g_2(\tilde{X}^{*i}(T^a)_i{}^j X_j \lambda^a + \lambda^{\dagger a} X^{\dagger i}(T^a)_i{}^j \tilde{X}_j - \tilde{X}^{*i}(T^a)_j{}^i \bar{X}_j \lambda^a - \lambda^{\dagger a} \bar{X}^{\dagger i}(T^a)_j{}^i \tilde{X}_j) \\ & + \frac{1}{2}(D^\mu \tilde{X}^* D_\mu \tilde{X} - M^2 |\tilde{X}|^2) - \frac{1}{2}M(\bar{X}_i X_i + X^{\dagger i} \bar{X}^{\dagger i}) + \frac{1}{2}g_2 D_a \tilde{X}^i (T^a)_i{}^j \tilde{X}_j \\ & - \frac{1}{2}M_{soft}^2 |\tilde{X}|^2.\end{aligned}\quad (1)$$

The component field notation has been used. In eq.(1), T^a 's are generators of the $SU(2)_L$ in **n** representation. X_i and \bar{X}^i consist of the left-hand pairs of X_i , transforming in $SU(2)_L$ **5** representation with the generator $(T^a)_i{}^j$ and the complex conjugate representation with the generator $(T^a)_i{}^j = (T^a)_j{}^i$, respectively. They are not independent. Actually they are dual to each other under the $SU(2)_L$. We write the Lagrangian in the form of eq.(1) just for convenience. Their superpartners compose a bosonic $SU(2)_L$ 5-plets \tilde{X}^i . Both X_i and \tilde{X}^i have trivial $SU(3)_c \times U(1)_Y$ quantum numbers (1, 0).

In terms of 4-component notation, we can define following spinors,

$$\Psi^{+2} \equiv \begin{pmatrix} X^{+2} \\ (\bar{X}^{+2})^\dagger \end{pmatrix}, \Psi^{-2} \equiv \begin{pmatrix} X^{-2} \\ (\bar{X}^{-2})^\dagger \end{pmatrix} = (\Psi^{+2})^C, \quad (2)$$

$$\Psi^{+1} \equiv \begin{pmatrix} X^{+1} \\ (\bar{X}^{+1})^\dagger \end{pmatrix}, \Psi^{-1} \equiv \begin{pmatrix} X^{-1} \\ (\bar{X}^{-1})^\dagger \end{pmatrix} = (\Psi^{+1})^C, \quad (3)$$

$$\Psi^0 \equiv \begin{pmatrix} X^0 \\ (\bar{X}^0)^\dagger \end{pmatrix} = (\Psi^0)^C. \quad (4)$$

The neutral field Ψ^0 is a Majorana field.

The superpotential takes the simple form:

$$W = W_{MSSM} + \frac{1}{2}M\tilde{X}^2, \quad (5)$$

which gives X and \tilde{X} the same unbroken supersymmetry mass M . \tilde{X} get a soft mass after supersymmetry breaking. The D -term contribution to the scalar potential is:

$$V_D = \frac{1}{2}g_2^2 \left[\sum \phi^* t^a \phi + \frac{1}{2} \tilde{X}^* T^a \tilde{X} \right]^2. \quad (6)$$

where ϕ denotes the $SU(2)_L$ scalars in the MSSM. Compared with the MSSM, the extra term is:

$$\frac{1}{2}g_2^2 \left[\sum (\phi^* t^a \phi)(\tilde{X}^* T^a \tilde{X}) + \left(\frac{1}{2} \tilde{X}^* T^a \tilde{X} \right)^2 \right]. \quad (7)$$

These couplings do not cause \tilde{X} decay but annihilation into MSSM $SU(2)_L$ scalars, which give an extra negligible mass splitting between \tilde{X}^i .

Considering non-renormalizable terms of the lagrangian, there are dimension 5 operators $\tilde{X}^{ijkl} \phi_i \phi_j \phi_k \phi_l / \Lambda$ for the complex scalar 5-plet \tilde{X} , and dimension 6 operators $X^{ijkl} \psi_i \phi_j \phi_k \phi_l / \Lambda^2$ for the fermionic 5-plet X allowed by the $SU(2)_L \times U(1)_Y$ gauge symmetry, where ψ_i is the left-hand leptons or the higgsinos in the MSSM, eg. $\tilde{X} H_u H_d H_u H_u^* / \Lambda$, $\tilde{X} H_u H_d H_d H_d^* / \Lambda$, $X \tilde{H}_u H_d H_d H_d^* / \Lambda^2$, $X L H_u H_d H_u / \Lambda^2$, etc.

We can generate these couplings by adding the corresponding higher dimension superpotential, eg.

$$W_{non-ren} = \frac{\tilde{X} H_u H_d H_u H_d}{\Lambda^2} + \frac{\tilde{X} \tilde{L} H_u H_d H_u}{\Lambda^2} + \dots, \quad (8)$$

the equation of motion for the auxiliary fields are:

$$\begin{aligned} F_{H_d} &= -\left(\frac{\partial W}{\partial H_d} \right)^* = -\left(\mu H_u + \frac{\tilde{X} H_u H_u H_d}{\Lambda^2} + \frac{\tilde{X} \tilde{L} H_u H_u}{\Lambda^2} + \dots \right)^*, \\ F_{H_u} &= -\left(\frac{\partial W}{\partial H_u} \right)^* = -\left(\mu H_d + \frac{\tilde{X} H_u H_d H_d}{\Lambda^2} + \frac{\tilde{X} \tilde{L} H_u H_d}{\Lambda^2} + \dots \right)^*. \end{aligned} \quad (9)$$

This generates dim6 couplings for the fermionic 5-plet : $X \tilde{H}_u H_d H_d H_d^* / \Lambda^2$, $X L H_u H_d H_u / \Lambda^2$ where $\Lambda \approx 10^{15} \text{GeV}$. These operators can induce 4-body decays with a typical life-time $\tau \sim \Lambda^4 \text{TeV}^{-5} \sim 10^{19} \text{s}$ which is longer than the age of the universe ($\sim 10^{17} \text{s}$). So these couplings have no influence on the observed stability of the DM candidates. The F-term also generates the scalar dim.5 operators but which are not suppressed by one power of Λ but two powers: $\mu \tilde{X} H_u H_d H_d H_d^* / \Lambda^2$, $\mu \tilde{X} \tilde{L} H_d H_d H_d^* / \Lambda^2$. The typical life-time of these operators is: $\tau \sim \Lambda^3 \text{TeV}^{-3} \mu^{-1} \sim 10^{20} \text{s}$ which is long enough. Of course there are even higher dimension couplings of the complex scalar \tilde{X} , eg. $\tilde{X} \tilde{X}^* H_u H_u^* H_d H_d^* H_d H_d^* / \Lambda^4$, $\tilde{X} \tilde{X}^* \tilde{L} H_u H_d H_u^* H_d^* / \Lambda^4$, etc. These higher dimension operators can be neglected in considering the decay.

Therefore the new introduced particles ($\tilde{X}^{\pm 2}, \tilde{X}^{\pm 1}, \tilde{X}^0$) only decay into ($X^{\pm 2}, X^{\pm 1}, X^0$) via gauge interactions. ($X^{\pm 2}, X^{\pm 1}, X^0$) are quite stable. We will further study mass splitting among them in the next section, and see that the DM candidate is still X^0 .

III. PROPERTIES OF THE SMDM

The DM candidate in the SMDM model is still X^0 . Mass splitting due to SUSY particles is small because of SUSY breaking.

A. Mass splitting

Mass splitting should be studied in detail, like that in the MDM, because it can be calculated with little uncertainties in this simple model. Because of EW symmetry breaking, the gauge kinetic terms gives the fermionic 5-plet X a mass splitting through loop corrections [1, 2], $\Delta M_{EW}^Q \equiv M^Q - M^0 \approx Q^2 \times 166 \text{ MeV}$, where M^Q and M^0 are the pole masses of X^Q and X^0 , respectively.

The scalar particles \tilde{X} 's also contribute to mass splitting of X . They are heavier than the fermions X 's by a soft mass M_{soft} which is generally expected to be about 100 GeV-1 TeV. This further mass splitting is calculated by using the supersymmetric kinetic term, the second line of the eq.(1), at the loop level which involves \tilde{X} 's and the gauginos. By using the two-component notation for fermions, the one-loop pole mass is written as [8]

$$M_{SUSY}^Q = M(1 + \frac{1}{2}\Sigma_L^Q + \frac{1}{2}\Sigma_R^Q) \quad (10)$$

where Σ_L^Q, Σ_R^Q are the 1PI self-energy functions as shown in Fig.1 in $Q = 0, +1$ cases.

In the diagrams of Fig.1, we denote the correspondent superpartners :

$$\phi^{+2} \equiv \tilde{X}^{+2}, \phi'^{-2} \equiv \tilde{X}^{+2}, \phi^{+1} \equiv \tilde{X}^{+1}, \phi'^{-1} \equiv \tilde{X}^{+1}, \phi^0 \equiv \tilde{X}^{+1}. \quad (11)$$

$\tilde{w}^{\pm}, \tilde{w}_3$ are the superpartners of the $SU(2)_L$ gauge bosons and $\tilde{w}^+ = V_{11}^{-1}\tilde{C}_1^+ + V_{12}^{-1}\tilde{C}_2^+$, $\tilde{w}^- = U_{11}^{-1}\tilde{C}_1^- + U_{12}^{-1}\tilde{C}_2^-$, $\tilde{w}_3 = N_{2i}^{-1}\tilde{N}_i$, $i=1-4$. $\tilde{C}_1^{\pm}, \tilde{C}_2^{\pm}$ and \tilde{N}_i are the charginos and neutralinos of the MSSM. U, V, N are the unitary matrices diagonalizing the mass matrices of charginos and neutralinos [5].

It is worthy to note that the Σ_D^Q term which is related to the B-term $B^{ij}\tilde{X}_i\tilde{X}_j$, may also appear in the pole mass formula, and it does not cause divergences. But in our calculation

$$\begin{aligned}
& \text{Diagram 1: } \text{Circle with } p \text{ and } -ip \cdot \bar{\sigma} \Sigma_L^{+1} = \text{Diagram 1.1} + \text{Diagram 1.2} + \text{Diagram 1.3} \\
& \text{Diagram 2: } \text{Circle with } p \text{ and } -ip \cdot \bar{\sigma} \Sigma_R^{+1} = \text{Diagram 2.1} + \text{Diagram 2.2} + \text{Diagram 2.3} \\
& \text{Diagram 3: } \text{Circle with } p \text{ and } -ip \cdot \bar{\sigma} \Sigma_L^0 = \text{Diagram 3.1} + \text{Diagram 3.2} \\
& \text{Diagram 4: } \text{Circle with } p \text{ and } -ip \cdot \bar{\sigma} \Sigma_R^0 = \text{Diagram 4.1} + \text{Diagram 4.2}
\end{aligned}$$

FIG. 1: One-loop corrections to the 1PI self-energy functions to the $Q = 0, +1$ components of the SMDM.

we do not consider it for simplicity, it is enough for us to break the supersymmetry only through the soft mass M_{soft} .

Using the superpartner notation mentioned in Sect. I we get :

$$\begin{aligned}
\Sigma_L^{+1} = & \frac{g_2^2}{16\pi^2} [V_{11}^* V_{11} B_1(\tilde{C}_1, \phi^{+2}) + V_{21}^* V_{21} B_1(\tilde{C}_2, \phi^{+2}) + \frac{3}{2} (U_{11}^* U_{11} B_1(\tilde{C}_1, \phi^0) \\
& + U_{21}^* U_{21} B_1(\tilde{C}_2, \phi^0)) + \frac{1}{2} (N_{12}^* N_{12} B_1(\tilde{N}_1, \phi^{+1}) + N_{22}^* N_{22} B_1(\tilde{N}_2, \phi^{+1}) \\
& + N_{32}^* N_{32} B_1(\tilde{N}_3, \phi^{+1}) + N_{42}^* N_{42} B_1(\tilde{N}_4, \phi^{+1})], \tag{12}
\end{aligned}$$

$$\begin{aligned}
\Sigma_R^{+1} = & \frac{g_2^2}{16\pi^2} [U_{11}^* U_{11} B_1(\tilde{C}_1, \phi'^{-2}) + U_{21}^* U_{21} B_1(\tilde{C}_2, \phi'^{-2}) + \frac{3}{2} (V_{11}^* V_{11} B_1(\tilde{C}_1, \phi^0) \\
& + V_{21}^* V_{21} B_1(\tilde{C}_2, \phi^0)) + \frac{1}{2} (N_{12}^* N_{12} B_1(\tilde{N}_1, \phi'^{-1}) + N_{22}^* N_{22} B_1(\tilde{N}_2, \phi'^{-1}) \\
& + N_{32}^* N_{32} B_1(\tilde{N}_3, \phi'^{-1}) + N_{42}^* N_{42} B_1(\tilde{N}_4, \phi'^{-1})], \tag{13}
\end{aligned}$$

$$\begin{aligned}\Sigma_L^0 = \frac{g_2^2}{16\pi^2} & \left[\frac{3}{2} (V_{11}^* V_{11} B_1(\tilde{C}_1, \phi^{+1}) + V_{21}^* V_{21} B_1(\tilde{C}_2, \phi^{+1}) + U_{11}^* U_{11} B_1(\tilde{C}_1, \phi^{-1}) \right. \\ & \left. + U_{21}^* U_{21} B_1(\tilde{C}_2, \phi^{-1})) \right],\end{aligned}\quad (14)$$

$$\begin{aligned}\Sigma_R^0 = \frac{g_2^2}{16\pi^2} & \left[\frac{3}{2} (U_{11}^* U_{11} B_1(\tilde{C}_1, \phi'^{-1}) + U_{21}^* U_{21} B_1(\tilde{C}_2, \phi'^{-1}) + V_{11}^* V_{11} B_1(\tilde{C}_1, \phi'^{+1}) \right. \\ & \left. + V_{21}^* V_{21} B_1(\tilde{C}_2, \phi'^{+1})) \right],\end{aligned}\quad (15)$$

where B_1 is the one rank two point integral

$$B_1(p^2, m_1, m_2) = -\frac{1}{2\varepsilon} + \frac{A_0(m_1) - A_0(m_2) + (m_2^2 - m_1^2 - p^2)B_0(p^2, m_1, m_2)}{2p^2} \quad (16)$$

with A_0 and B_0 being the Passarino-Veltman functions.

Because all the superpartners ϕ^i have the same mass $M + M_{soft}$, we can simplify the above four equations to get the final result of the mass splitting due to SUSY particles :

$$\begin{aligned}\Delta M_{SUSY}^Q &= \frac{Q^2}{2} (\Sigma_L^Q + \Sigma_R^Q - \Sigma_L^0 - \Sigma_R^0) \\ &= \frac{g_2^2 Q^2}{16\pi^2} \left[-(V_{11}^* V_{11} + U_{11}^* U_{11}) B_1(\tilde{C}_1, \phi) - (V_{21}^* V_{21} + U_{21}^* U_{21}) B_1(\tilde{C}_2, \phi) \right. \\ &\quad \left. + \frac{1}{2} N_{i2}^* N_{i2} B_1(\tilde{N}_i, \phi) \right].\end{aligned}\quad (17)$$

The poles in the B_1 function are canceled as expected using the unitarity of the U, V and N . The mass splitting is a function of $M_1, M_2, \tan \beta, \mu, M_{soft}$ and M . In the correct EW breaking parameter space, our numerical result for the mass splitting due to SUSY particles is that

$$\Delta M_{SUSY}^Q \sim 0.01 Q^2 \text{ MeV} \quad (18)$$

which is negligibly small compared to the pure EW corrections.

B. The thermal relic density

The thermal relic density fixes the WIMP mass. In the MDM, the relic species are $\{X^{\pm 2}, X^{\pm 1}, X^0\}$ and the coannihilation channels are:

$$X^i X^j \rightarrow AA, f \bar{f}. \quad (19)$$

where A and f denote a EW gauge boson and the SM fermion, respectively. The mass splittings among them are very small compared to their masses. In the density calculation,

such mass splittings are negligible. The relic particle thermal average cross section is[1]:

$$\langle\sigma_{Av}\rangle(X^i X^j \rightarrow AA, f\bar{f}) \approx \frac{\pi\alpha_2^2}{8M^2} \times 166, \quad (20)$$

Matching to the relic abundance, the DM particle mass is determined to be $M = 4.4$ TeV without considering Sommerfeld corrections.

Once SUSY is introduced in, a whole bunch of superpartners of those in MDM will present. SUSY breaking gives soft masses to the scalars and gauginos. Looking at \tilde{X} 's, they have a mass $M + M_{soft}$. In general when $M_{soft} \leq 0.1M$, eg. $M_{soft} \leq 440$ GeV for $M = 4.4$ TeV, they have sizable effects on the relic abundance and must be included in the the relic species [6, 7]. So now the coannihilation relic species are $\{(\tilde{X}^{\pm 2}, \tilde{X}^{\pm 1}, \tilde{X}^0), (X^{\pm 2}, X^{\pm 1}, X^0)\}$ and the coannihilation channels are:

$$\begin{aligned} \tilde{X}^i \tilde{X}^j &\rightarrow AA, f\bar{f}, \tilde{G}\tilde{G}, \\ \tilde{X}^i X^j &\rightarrow f\tilde{f}, \tilde{G}A, \\ X^i X^j &\rightarrow AA, f\bar{f}, \tilde{G}\tilde{G}, \end{aligned} \quad (21)$$

where \tilde{G} denotes MSSM gauginos and \tilde{f} the superpatner of f .

Furthermore, because M is also much larger than the electroweak scale, the physics of the above-mentioned coannihilation is basically supersymmetric and EW gauge symmetric, we can make unbroken EW symmetry and unbroken supersymmetry approximation when calculating the thermal average cross section. Nevertheless, it is still a hard work. In terms of two-component fields, there are 24 gauge kinetic vertices and another 24 vertices involving superpatners. But actually we can have a useful and proper estimate for the relations between cross sections for the three kinds of processes in eq. (21). It is found that introducing in SUSY have nearly 4 times influence on the thermal average cross section. It will be shown in Fig.2-8 a series of explicit examples by using two-component spinor techniques[8] and the results reads:

$$v\sigma(\bar{X}^{+2} X^{+2} \rightarrow W_3 W_3) = 16 \times \frac{8\pi\alpha^2}{3M^2}, \quad (22)$$

$$v\sigma(\phi^{*+2} \phi^{+2} \rightarrow W_3 W_3) = 32 \times \frac{8\pi\alpha^2}{3M^2}, \quad (23)$$

$$v\sigma(\bar{X}^{+2} X^{+2} \rightarrow \tilde{\omega}_3 \tilde{\omega}_3) \sim 0, \quad (24)$$

$$v\sigma(\phi^{*+2} \phi^{+2} \rightarrow \tilde{\omega}_3 \tilde{\omega}_3) \sim 0, \quad (25)$$

$$v\sigma(\phi^{+2}\phi'^{-2} \rightarrow \tilde{\omega}_3\tilde{\omega}_3) \simeq 16 \times \frac{3\pi\alpha^2}{2M^2}, \quad (26)$$

$$v\sigma(\bar{X}^{+2}\phi^{+2} \rightarrow W_3\tilde{\omega}_3) \simeq 16 \times \frac{3\pi\alpha^2}{8M^2}, \quad (27)$$

$$v\sigma(X^{+2}\phi'^{-2} \rightarrow W_3\tilde{\omega}_3) \simeq 16 \times \frac{3\pi\alpha^2}{8M^2}, \quad (28)$$

where v is the relative velocity in the lab frame, Eq. (24),(25) is the results of the p-wave suppression.




FIG. 2: The eight Feynman diagrams for $\bar{X}^{+2}X^{+2} \rightarrow W_3W_3$




FIG. 3: The three Feynman diagrams for $\phi^{*+2}\phi^{+2} \rightarrow W_3W_3$




FIG. 4: The four Feynman diagrams for $\bar{X}^{+2}X^{+2} \rightarrow \tilde{\omega}_3\tilde{\omega}_3$




FIG. 5: The two Feynman diagrams for $\phi^{*+2}\phi^{+2} \rightarrow \tilde{\omega}_3\tilde{\omega}_3$




FIG. 6: The four Feynman diagrams for $\phi^{+2}\phi'^{-2} \rightarrow \tilde{\omega}_3\tilde{\omega}_3$




FIG. 7: The twelve Feynman diagrams for $\bar{X}^{+2}\phi^{+2} \rightarrow W_3\tilde{\omega}_3$ and $X^{+2}\phi'^{-2} \rightarrow W_3\tilde{\omega}_3$

We have already calculate a complete series of thermal cross section, we end up with:

$$v\sigma_{eff} = \sum v\sigma \approx 4v\sigma(\bar{X}^{+2}X^{+2} \rightarrow W_3W_3) \quad (29)$$

with considering the freedom for X^{+2} , \bar{X}^{+2} and ϕ^{+2}, ϕ'^{-2} are all $g = 2$. This is very different with the dramatically influence from the higher representation such as $SU(2)$ -5 in our problem. Despite we make this conclusion from some specific examples, we think it is a common

result.

Once the DM particle is as heavy as few TeV, the Sommerfeld effect due to hundreds GeV particles should be taken into consideration. In the MDM, the Sommerfeld enhancement effect due to SM particles, especially due to W and Z bosons, has been calculated, the factor is about 5 [9]. We expect approximately the same result in our case. While our case is $N = 1$ SUSY, all the SUSY particles should be included in the ladder diagram calculation in determining the potential which tells us the enhancement factor. However including the SUSY particles does not change the Sommerfeld effect essentially. In the extreme case of $N = 4$ SUSY, the extra symmetries just keep the gauge coupling from running. The potential itself has the same form as in the non-SUSY Yang-Mills case. In our $N = 1$ SUSY case with soft breaking, the logarithmic running of the gauge coupling of the MDM is expected only mildly reduced, the Sommerfeld effect is then approximately the same. So it is reasonable to say that the Sommerfeld enhancement factor is about the same as that calculated in the MDM, $M \simeq \sqrt{5} \times 2 \times 4.4 \simeq 19.7$ TeV.

C. Direct and Indirect signatures

As for the direct DM detection rate, this SMDM is of the same order as the MDM. The DM particle interacts with quarks via loops. Although more heavy particles appear in the loops, the total cross section has the same order, $\sigma_{\text{SI}} \propto 10^{-44} \text{cm}^2$, which is within the sensitive of the current experiments, such as Super-CDMS and Xenon 1-ton[10, 11].

The DM annihilation in the galaxy may have observable signatures. The estimation of the cross section is also like that in the MDM[12, 13], the result has no order change. Note that SUSY does not change the Sommerfeld effect essentially, the predominant annihilation channel is still into EW W bosons, $\langle \sigma v \rangle_{WW} \sim 10^{-23} \text{cm}^3 \text{s}^{-1}$. Because the DM mass $M \simeq 19.7$ TeV, we expect this model predict: (1) continuous rise $e^+/e^+ + e^-$ spectrum up to about 20 TeV; (2) flat $e^+ + e^-$ spectrum up to M ; (3) \bar{p}/p flux has excess above the energy probed by PAMELA. SMDM has the solutions to the PAMELA and Fermi-LAT anomaly[14, 15], the predictions in higher energies need further experimental data to verify.

IV. SUMMARY AND DISCUSSION

We have made SUSY extension to the MDM model by introducing a complex scalar quintuplet as the superpartner of the fermion quintuplet. The neutral component of the fermionic 5-plet is still the DM particle as in the MDM model. Mass splittings among the fermionic 5-plets due to SUSY have been calculated in detail, they are found to be small. By considering new relic species and new coannihilation channels into the MSSM final states, the DM mass is estimated to be 19.7 TeV.

The direct and indirect signals are basically the same as those in the MDM. Numerically the DM elastic scattering cross section with a nucleus is about $10^{-44} \text{ cm}^3 \text{ s}^{-1}$ and the cross section of the predominant annihilation channel into W bosons is about $10^{-23} \text{ cm}^3 \text{ s}^{-1}$. SMDM predicts $e^+, e^+ + e^-, \bar{p}$ spectrum in agreement with the previous PAMELA, Fermi-LAT data, \bar{p} flux has excess above the energy probed by PAMELA which need further experimental test.

In the near future, suppose SUSY is discovered, say at LHC, it will be still a question if MSSM itself provides a DM particle, because R-parity as a discrete symmetry is still an assumption which is not as solid as gauge invariance and SUSY. It is plausible that R-parity is violated. In that case, it is still simple and interesting to have the DM via introducing $SU(2)_L$ high dimensional representations.

Acknowledgments

We would like to thank Yu-Qi Chen, Xiao-Jun Bi, Jia-Shu Lu and Hua Shao for helpful discussions. This work was supported in part by the National Natural Science Foundation of China under Nos. 11075193 and 10821504, and by the National Basic Research Program of China under Grant No. 2010CB833000.

Appendix: the su(2)-n representation

The SU(2)-n representation U^j ($n=2j+1$) is self-conjugate , when j is integer (n -odd), U^j is real , when j is half-integer (n -even), U^j is self-conjugate also but not real :

$$XU^jX^{-1} = U^{j*} \Rightarrow \begin{cases} U^j = U^{j*} & j = 0, 1, 2, \dots, X \text{ is symmetric} \\ U^j \text{ is not real} & j = \frac{1}{2}, \frac{3}{2}, \dots, X \text{ is antisymmetric} \end{cases} \quad (\text{A.1})$$

it is important in proving some identities including the SU(2) generators.

The generators of the su(2)-n representation is:

$$\begin{aligned} (T_1^j)_{\nu\mu} &= \frac{1}{2}[\delta_{\nu(\mu+1)}\Gamma_\nu^j + \delta_{\nu(\mu-1)}\Gamma_{-\nu}^j] \\ (T_2^j)_{\nu\mu} &= -\frac{i}{2}[\delta_{\nu(\mu+1)}\Gamma_\nu^j - \delta_{\nu(\mu-1)}\Gamma_{-\nu}^j] \\ (T_3^j)_{\nu\mu} &= \mu\delta_{\nu\mu}. \end{aligned} \quad (\text{A.2})$$

where

$$\Gamma_\nu^j = \Gamma_{-\nu+1}^j = (j + \nu)(j - \nu + 1)^{1/2} \quad (\text{A.3})$$

for example

$$T_3^1 = \frac{\sqrt{2}}{2} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, T_3^2 = \frac{\sqrt{2}}{2} \begin{pmatrix} 0 & -i & 0 \\ i & 0 & -i \\ 0 & i & 0 \end{pmatrix}, T_3^3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad (\text{A.4})$$

$$T_5^1 = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & \frac{\sqrt{6}}{2} & 0 & 0 \\ 0 & \frac{\sqrt{6}}{2} & 0 & \frac{\sqrt{6}}{2} & 0 \\ 0 & 0 & \frac{\sqrt{6}}{2} & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix}, T_5^2 = \begin{pmatrix} 0 & -i & 0 & 0 & 0 \\ i & 0 & -i\frac{\sqrt{6}}{2} & 0 & 0 \\ 0 & i\frac{\sqrt{6}}{2} & 0 & -i\frac{\sqrt{6}}{2} & 0 \\ 0 & 0 & i\frac{\sqrt{6}}{2} & 0 & -i \\ 0 & 0 & 0 & i & 0 \end{pmatrix}, T_5^3 = \begin{pmatrix} 2 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & -2 \end{pmatrix}. \quad (\text{A.5})$$

and have

$$\bar{T}^a T^2 = -T^2 T^a \quad (\text{A.6})$$

[1] M. Cirelli, N. Fornengo, A. Strumia, Nucl.Phys. B753 (2006) arXiv:hep-ph/0512090.

- [2] M. Cirelli and A. Strumia, New J. Phys. 11 (2009) 105005 arXiv:0903.3381.
- [3] Y. Cai, X.-G. He, M. Ramsey-Musolf, and L.-H. Tsai, arXiv:1108.0969;
C.-H. Chen and S.S.C. Law, arXiv:1111.5462.
- [4] For reviews, see M. E. Peskin, arXiv:0801.1928; S. P. Martin, arXiv:hep-ph/9709356.
- [5] G. Bertone, D. Hooper and J. Silk, Phys. Rep. **405**, 279-390, (2005).
- [6] G. Jungman and M. Kamionkowski and K. Griest, Phys. Rep. **267**, 195 (1996).
- [7] J. Edsjo and P. Gondolo, Phys. Rev. D **56**,1879-1894,(1997).
- [8] H. K. Dreiner, H. E. Haber and S. P. Martin, arXiv:0812.1594.
- [9] M. Cirelli, A. Strumia, Matteo Tamburini, Nucl.Phys. B787 (2007), arXiv:0706.4071.
- [10] Z.Ahmed *et al.* [CDMS Collaboration], Phys. Rev. Lett. 102, 011301 (2009).
- [11] J.Angle *et al.* [XENON Collaboration], arXiv: 1104.2549.
- [12] M. Cirelli and A. Strumia, PoS IDM 2008 (2008) 089 arXiv: 0808.3867.
- [13] M. Cirelli, R. Franceschini, A. Strumia Nucl. Phys. B 800:204-220,2008 arXiv:0802.3378.
- [14] O. Adriani *et al.* [PAMELA Collaboration], arXiv:1103.2880.
- [15] A. A. Abdo *et al.* [Fermi LAT Collaboration], Phys. Rev. Lett. 102 (2009).